arXiv:1210.4422v1 [cond-mat.mes-hall] 16 Oct 2012

Magnetically robust topological edge states and flat bands

Tomi Paananen and Thomas Dahm[1]

[1] *Universität Bielefeld, Fakultät für Physik, Postfach 100131, D-33501 Bielefeld, Germany*

(Dated: October 17, 2012)

We study thin strips of three dimensional topological insulators in the presence of a spin-splitting Zeeman field. We show that under certain conditions the topological edge states at the sides of a strip remain robust against a time-reversal symmetry breaking Zeeman field. For a particle-hole symmetric system with Zeeman field perpendicular to the strip we strictly proof that the dispersion and the spin-orbital structure of the edge states remains unchanged. When the Zeeman field lies parallel to the strip, the Dirac spectrum becomes flat, but remains intact. Above a critical value of the Zeeman field a completely flat surface band develops. We present numerical calculations for a realistic tight-binding model of Bi_2Se_3. These calculations show that even though particle-hole symmetry is not strictly fulfilled in this system, these special features are still present.

PACS numbers: 73.20.At, 75.70.-i, 73.43.-f, 03.65.Vf

Symmetries play an important role in condensed matter physics. Recently, a peculiar state of matter, the topological insulator, has been first suggested theoretically [1, 2] and subsequently observed experimentally [3–8]. In these materials time-reversal symmetry guarantees protection of conducting surface states against backscattering, which is a requirement for dissipationless spin transport. Usually, when a time-reversal symmetry breaking perturbation is applied, like a magnetic field for example, the surface states are not protected anymore. Here, we will show that under certain circumstances topological surface states may become robust even under a time-reversal symmetry breaking perturbation. Also, a robust flat band may appear.

Topological insulators are materials, which have an insulating energy gap in their bulk, but possess topologically protected gapless surface states due to significant spin-orbit coupling. Due to Kramers theorem a time-reversal invariant perturbation cannot open a gap in the Dirac spectrum of the surface states in these systems. However, a gap can be opened, when a time-reversal breaking perturbation is applied.

Recently, several materials, like Bi_2Se_3, Bi_2Te_3, Sb_2Te_3, or $PbBi_2Te_4$ have been experimentally identified as three dimensional topological insulators [5–8]. The topological protection of the surface states and their spin-momentum coupled nature makes these systems particularly interesting for spintronics. Several devices have been proposed theoretically [9–13]. In particular, the combination with ferromagnetic materials or the application of magnetic fields can lead to interesting effects.

In the present work we study thin strips of three dimensional topological insulators in the presence of a time-reversal symmetry breaking Zeeman field. Zeeman fields can be introduced into topological insulators either by doping with ferromagnetic dopants [13] or by proximity to a ferromagnetic material. We show that in this case certain additional spin-orbital symmetries are present, which are respected by the Zeeman field. These symmetries can make the surface states robust even against a Zeeman field. We will also show that at larger Zeeman fields a phase transition appears into a state with a robust flat band. Flat bands in solids are particularly interesting objects, because the group velocity vanishes, allowing for immobile wave packets, localized states, and a giant effective mass [18].

We start from the generic effective two-orbital tight-binding model for a three dimensional topological insulator suggested by Li et al [15, 17] in the presence of a Zeeman field:

$$H(\mathbf{k}) = \epsilon_0(\mathbf{k})\mathbb{I}_{4\times4} + \sum_{i=0}^{2} m_i(\mathbf{k})\Gamma^i + \sum_{\alpha\in\{x,y,z\}} V_\alpha\, \mathbb{I}_{2\times2}\otimes\sigma_\alpha \tag{1}$$

with $\epsilon_0(\mathbf{k}) = C+2D(1-\cos k_x)+2D(1-\cos k_y)$, $m_0(\mathbf{k}) = M - 2B(1-\cos k_x) - 2B(1-\cos k_y)$, $m_1(\mathbf{k}) = 2A\sin k_x$, and $m_2(\mathbf{k}) = 2A\sin k_y$. Here, the Dirac Γ matrices are represented by $\Gamma^{0,1,2} = (\tau_z\otimes\mathbb{I}_{2\times2}, \tau_x\otimes\sigma_x, \tau_x\otimes\sigma_y)$ in the basis of bonding and antibonding p_z orbital states. The Pauli matrices in orbital space are denoted by τ_i and the ones in spin space by σ_i. The components of the Zeeman field in x, y, and z-direction are denoted by $V_{x,y,z}$, respectively. The parameters A, B, C, D, and M have been derived from bandstructure calculations for Bi_2Se_3 in Ref. 17.

First, we will study a particle-hole symmetric system with $D = 0$. In Bi_2Se_3 the parameter D is about a factor of three smaller than B and its modest influence will be considered further below. Without loss of generality we can set $C = 0$. The Hamiltonian (1) then possesses several spin-orbital symmetries in the absence of a Zeeman field some of which remain valid for certain directions of the Zeeman field. Specifically, let us consider the following symmetry operators: $\Theta_0 = \tau_z\otimes\sigma_x$, $\Theta_1 = \tau_x\otimes\sigma_x$, $\Theta_2 = \tau_y\otimes\sigma_y$, $\Theta_3 = \tau_z\otimes\sigma_z$, and parity $P = \tau_z\otimes\mathbb{I}_{2\times2}$. In the absence of a Zeeman field the Hamiltonian (1) fulfils the following symmetry relations: $\Theta_0 H(k_x,k_y)\Theta_0^{-1} = H(-k_x,k_y)$,

$\Theta_1 H(k_x,k_y)\Theta_1^{-1} = -H(-k_x,k_y)$, $\Theta_2 H(k_x,k_y)\Theta_2^{-1} = -H(-k_x,k_y)$, $\Theta_3 H(k_x,k_y)\Theta_3^{-1} = H(k_x,k_y)$, and $PH(k_x,k_y)P^{-1} = H(-k_x,-k_y)$. In the presence of a Zeeman field in z-direction the symmetries Θ_1, Θ_2, Θ_3, and P are still respected, while Θ_0 is broken. In contrast, in the presence of a Zeeman field in x-direction the symmetries Θ_0, Θ_2, and P are still respected, while Θ_1 and Θ_3 are broken. All of these symmetry operations respect $\Theta_i^2 = P^2 = \mathbb{I}_{4\times 4}$, thus their eigenvalues are ± 1.

Let us consider now a strip with finite extension in y-direction with $y \in [0, L]$, but periodic boundary conditions in x-direction. In the absence of a Zeeman field the system possesses four topological surface states at the two edges $y = 0$ and $y = L$, two on each side. These edge states can be obtained analytically in small k_y expansion following the method of Ref. 19. If L is sufficiently larger than the localization length of the edgestates, we may consider the edges separately letting $L \to \infty$. The edge state wavefunctions are then linear superpositions of exponentially decaying functions of the form $e^{\eta y}$ with the boundary conditions $\psi(y = 0) = 0$ and $\psi(y \to \infty) = 0$.

We will first proof that the dispersion of the edge states remains unaffected by a Zeeman field in z-direction. Consider the Hamiltonian

$$H_0(\mathbf{k}) = m_0(\mathbf{k})\Gamma^0 + m_2(\mathbf{k})\Gamma^2 + V_z\, \mathbb{I}_{2\times 2} \otimes \sigma_z \qquad (2)$$

This is the Hamiltonian (1) with $C = D = 0$ except for the Γ^1 term. Note, that this Hamiltonian anticommutes with the symmetry operation Θ_1, i.e. $[H_0, \Theta_1]_+ = 0$. Now assume that $|\Psi_{k_x}\rangle$ is a zero-energy eigenstate of H_0. It is easy to show by direct calculation that such a zero-energy edge state exists, if $M - 2B(1 - \cos k_x) + V_z > 0$. A second one exists, if $M - 2B(1 - \cos k_x) - V_z > 0$. Physically, this means that an edge state exist for each of the two spin-split bulk bands whenever there is a band inversion [1]. Due to the anticommutation, $|\Psi_{k_x}\rangle$ either is an eigenstate of Θ_1 or can be constructed as such. Now it turns out that $\Theta_1 = \Gamma^1$. Therefore $|\Psi_{k_x}\rangle$ will also be an eigenstate of the Hamiltonian (1) (with $C = D = 0$) and its energy dispersion will thus be given by the prefactor of the Γ^1 term times the Θ_1 eigenvalue, i.e. $E(k_x) = \pm 2A \sin k_x$. This result is apparently independent of V_z (and also of M and B, as long as the condition for the existence of the edge state is fulfilled). As the Hamiltonian also commutes with Θ_3 we can deduce that the four edge states have to be proportional to $(1, 0, 0, 1)$, $(1, 0, 0, -1)$, $(0, 1, 1, 0)$, or $(0, 1, -1, 0)$, i.e. also the spin-orbital structure of the edge states remains unaffected by V_z. It is only the spatial structure that changes. Note, that this spin-orbital structure has the property that the expectation value of the total spin-z component vanishes, because the spin in the two orbital subbands is just opposite. This is in contrast to the quantum spin Hall system HgTe, where the edge states consist of two orbital subbands having the same spin [19, 20]. In the present case, which is relevant for the edge states in a thin $\mathrm{Bi_2Se_3}$ strip, this special feature explains why the edge states are robust against a Zeeman field in z-direction.



FIG. 1: (Color online) Numerical dispersions of bulk and edge states for $V_z/M = 0.5$, 1.0, and 1.5, respectively. When V_z/M reaches 1 two of the four edge states disappear. We have used $B = A = M = 1$, and $C = D = 0$.

The parity operator P relates a state at one edge to a state with same energy and opposite k_x on the other edge. From this we can conclude that the states $(1, 0, 0, 1)$ and $(1, 0, 0, -1)$ are localized on different edges and have the same spatial structure and the states $(0, 1, 1, 0)$ and $(0, 1, -1, 0)$, too. For $V_z = 0$ the symmetry $\Theta_0\Theta_2$ shows that $(1, 0, 0, 1)$ and $(0, 1, -1, 0)$ are on the same edge with same momentum, but opposite energy. The same is true for $(1, 0, 0, -1)$ and $(0, 1, 1, 0)$. These statements also hold for finite V_z even though Θ_0 is broken.

In order to confirm our analytical results we have diagonalized the tight-binding Hamiltonian (1) with $C = D = 0$ and $M = B = A = 1$ on a finite lattice of size 500×200 numerically with periodical boundary conditions in x-direction. The numerical results, shown in Fig. 1 confirm the analytical proof. For $V_z/M < 1$ we find four edge states, two on each side. When V_z/M approaches 1, the localization length of two of the edge states approaches the width of the system, similar to what has been found in Ref. 21. At $V_z/M = 1$ the bulk bands touch the Dirac point (see Fig. 1(b)). When V_z/M becomes larger than 1, these two edge states disappear. The other two edge states survive, though. The system thus makes a quantum phase transition into a quantum Hall state. Throughout this transition the dispersion and the spin-orbital structure of the edge states remains unchanged, while the bulk spectrum changes, of course.

Let us next consider a Zeeman field in x-direction. In this case the dispersion of the edge states changes, because $\mathbb{I}_{2\times 2} \otimes \sigma_x$ does not anticommute with Θ_1 anymore. The symmetries Θ_0 and Θ_2 are still respected. Let $|\Psi_{k_x}\rangle$ be an edge state of H with energy E. Then, $\Theta_0\Theta_2\, |\Psi_{k_x}\rangle$ is an edge state with energy $-E$ which is localized at the same edge. If we have just one edge state on each side, $|\Psi_{k_x}\rangle = \pm\Theta_0\Theta_2\, |\Psi_{k_x}\rangle$ must hold and its energy thus has to be strictly zero, i.e. a flat band appears. As shown in Fig. 2, this happens for $V_x > \sqrt{M(k_x)^2 + 4A^2 \sin^2(k_x)}$, where $M(k_x) = M - 2B(1 - \cos k_x)$. This is a topological flat band as has been discussed recently in graphene [22] and topological superconductors with gap nodes [23].



FIG. 2: (Color online) Numerical dispersions of bulk and edge states for a Zeeman field in x-direction with $V_x/M = 0.6$, and 1.6, respectively. When V_x/M exceeds 1 two of the four edge states disappear and a flat band appears. Parameters are the same as in figure 1.



FIG. 3: (Color online) Numerical dispersions of bulk and edge states for Zeeman field in z-direction with $V_z/M = 0.5$, 1.0 and 1.5, respectively. When V_z/M exceeds 1, two of the four edge states disappear. Here, we have chosen realistic parameters for Bi_2Se_3, as described in the text.

Such a flat band appears when the bulk gap closes [24], as is the case here, too. In the present case the flat band is generated by a magnetic exchange field, however, and thus could be tuned experimentally. We will show below that it can be observed realistically in Bi_2Se_3.

In Fig. 2 we show the dispersions of bulk and edge states above and below $V_x/M = 1$. For $V_x/M < 1$ we have four edge states. When $V_x/M > 1$ two of these states disappear and a pair of bulk Dirac points appears. These points are connected by a flat band of two edge states, one on each side of the strip.

So far we have discussed the particle-hole symmetric case $C = D = 0$ in Eq. (1), in which the two phenomena of stable dispersion and flat bands become particularly clear. We will now study parameters, which are realistic for Bi_2Se_3, in order to show that these phenomena remain and how they are modified. According to Zhang et al. [17] the parameters are given by $B = 3.3\,\mathrm{eV}$, $M = 0.28\,\mathrm{eV}$, $A = 0.49\,\mathrm{eV}$, $D = 1.14\,\mathrm{eV}$, and $C = 0.0068\,\mathrm{eV}$, i.e. particle-hole symmetry is somewhat broken. The parameter M is the size of the bulk gap in the absence of a Zeeman field. The parameter $t = D/B = 0.35$ is a measure of the particle-hole asymmetry. For $t > 1$ the system would become topologically trivial.

Following the method of Zhou et al. [19] we can determine the dispersion and structure of the edge states analytically, when a Zeeman field in z-direction is applied. For the dispersions we find $E(k_x) = t(M \pm V_z) \pm 2A\sqrt{1-t^2}\sin k_x$. The spin-orbital structure of the edge states turns out to be of the form $(u, 0, 0, v)$, $(-u, 0, 0, -v)$, $(0, v, u, 0)$, and $(0, -v, u, 0)$, where $u = \sqrt{1-t^2}$ and $v = \sqrt{1+t^2}$. For $t = 0$ the particle-hole symmetric results above are reproduced. Interestingly, the dispersions remain robust except for a constant energy shift of $\pm tV_z$, which is reduced by the particle-hole asymmetry. The Zeeman field does not create a gap in the dispersion of the edge states. Also, the spin-orbital structure of the edge states neither depends on V_z nor on k_x. Only the parameter t determines the spin-orbital structure.



FIG. 4: (Color online) (a) Numerical dispersion of bulk and edge states for a Zeeman field in x-direction with $V_x/M = 2.2$. When V_x/M exceeds 1, two of the four edge states disappear and an almost flat band appears. (b) Density of States of this system for $V_x/M = 4$. The flat band appears as a peak around $E = 0.45\,\mathrm{eV}$. Parameters are the same as in figure 3.

In Fig. 3 we present numerical dispersions for a finite Bi_2Se_3 strip in the presence of a Zeeman field in z-direction using the parameters of Zhang et al, which corroborate our analytical results. Fig. 3(a) shows that V_z shifts the energies of the edge states, but does not change the shape of the dispersions. When V_z/M exceeds 1, two of the four edge states disappear and the system makes a quantum phase transition into a quantum Hall state.

When the Zeeman field is applied in x-direction, the symmetries Θ_0 and P are still respected, while Θ_2 is broken. Let $|\Psi_{k_x}\rangle$ be an edge state of H with energy E. Then, $\Theta_0 |\Psi_{k_x}\rangle$ is an edge state with energy E and momentum $-k_x$, which is localized at the same edge. If there exists only one state at each edge we have $E(k_x) = E(-k_x)$, i.e. the dispersion is symmetric in k_x. The state $P\Theta_0 |\Psi_{k_x}\rangle$ then is an edge state with same energy and momentum k_x localized at the opposite edge and the edge state dispersion is twofold degenerate.

In Fig. 4(a) we show the numerical dispersion of bulk and edge states for a Zeeman field in x-direction with $V_x/M = 2.2$. When V_x/M exceeds 1, again two of the four edge states disappear and an almost flat band appears. Due to the broken particle-hole symmetry this band has a finite dispersion now. For $V_x/M = 2.2$ this band is clearly separated from the bulk bands. It ends at the two Dirac points of the bulk bands, as in the particle-hole symmetric case. In Fig. 4(b) we show the density of states of the system for $V_x/M = 4$. Here, the flat band appears as a peak around $E = 0.45$ eV. In order to realize this situation experimentally one would need Zeeman fields in the range $0.28\,\mathrm{eV} < V_x < 1\,\mathrm{eV}$, which are typical exchange splittings in common ferromagnets. The criterion for the appearance of this flat band remains the same as in the particle-hole symmetric case above, i.e. $V_x > \sqrt{M(k_x)^2 + 4A^2\sin^2(k_x)}$, because its appearance is related to the closure of the bulk gap, which is unchanged, when $D > 0$. The dispersion of the band decreases with increasing value of V_x.

In conclusion, we have shown that spin-orbital symmetries in a topological insulator can create special features like robust edge states and flat bands. Specifically, we have studied a thin $\mathrm{Bi_2Se_3}$ strip in a Zeeman field. When the Zeeman field is applied perpendicular to the strip we have shown that the edge states remain robust even under the time-reversal symmetry breaking Zeeman field. Above a critical value the system can be brought into a quantum Hall state via a quantum phase transition. When the Zeeman field is applied parallel to the strip along its long direction, a topological flat band appears above a critical value of the Zeeman field. The combination of $\mathrm{Bi_2Se_3}$ thin films with ferromagnetic materials can thus create highly interesting features that can be tuned by the polarization of the ferromagnet and lead to new spintronics devices.

[1] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, Science **314**, 1757 (2006).
[2] L. Fu, C.L. Kane and E.J. Mele, Phys. Rev. Lett. **98**, 106803 (2007).
[3] M. König, S. Wiedmann, C. Brüne, A. Roth, H. Buhmann, L.W. Molenkamp, X.-L. Qi, and S.-C. Zhang, Science **318**, 766 (2007).
[4] D. Hsieh, D. Qian, L. Wray, Y. Xia, Y. Hor, R.J. Cava, and M.Z. Hasan, Nature (London) **452**, 970 (2008)
[5] Y.L. Chen, J.G. Analytis, J.-H. Chu, Z.K. Liu, S.-K. Mo, X.L. Qi, H.J. Zhang, D.H. Lu, X. Dai, Z. Fang, S.C. Zhang, I.R. Fisher, Z. Hussain, and Z.-X. Shen, Science **325**, 178 (2009).
[6] Y. Xia, D. Qian, D. Hsieh, L. Wray, A. Pal, H. Lin, A. Bansil, D. Grauer, Y.S. Hor, R.J. Cava, and M.Z. Hasan, Nature Phys. **5**, 398 (2009).
[7] D. Hsieh, Y. Xia, D. Qian, L. Wray, F. Meier, J.H. Dil, J. Osterwalder, L. Patthey, A.V. Fedorov, H. Lin, A. Bansil, D. Grauer, Y.S. Hor, R.J. Cava, and M.Z. Hasan, Phys. Rev. Lett. **103**, 146401 (2009).
[8] K. Kuroda et al, Phys. Rev. Lett. **108**, 206803 (2012).
[9] I. Garate and M. Franz, Phys. Rev. Lett. **104**, 146802 (2010).
[10] T. Yokoyama , Y. Tanaka, and N. Nagaosa, Phys. Rev. B **81**, 121401(R) (2010).
[11] P. Hosur, P. Ghaemi, R.S.K. Mong, and A. Vishwanath, Phys. Rev. Lett. **107**, 097001 (2011).
[12] A.M. Black-Schaffer and J. Linder, Phys. Rev. B **84**, 180509(R) (2011).
[13] R. Yu, W. Zhang, H.-J. Zhang, S.-C. Zhang, X. Dai, and Z. Fang, Science **329**, 61 (2010).
[14] Yi Zhang *et al.*, Nature Phys. **6**, 584 (2010).
[15] R. Li, J. Wang, X.-L. Qi, and S.-C. Zhang, Nature Phys. **6**, 284 (2010).
[16] R.-L. Chu, J. Shi, and S.-Q. Shen, Phys. Rev. B **84**, 085312 (2011).
[17] H. Zhang, C.-X. Liu, X.-L. Qi, X. Dai, Z. Fang, and S.-C. Zhang, Nature Phys. **5**, 438 (2009).
[18] S. Nishino and M. Goda, J. Phys. Soc. Japan **74**, 393 (2005).
[19] B. Zhou, H.-Z. Lu, R.-L. Chu, S.-Q. Shen, and Q. Niu, Phys. Rev. Lett. **101**, 246807 (2008).
[20] L. Hao and T.K. Lee, Phys. Rev. B **83**, 134516 (2011).
[21] M. Wada, S. Murakami, F. Freimuth, and G. Bihlmayer, Phys. Rev. B **83**, 121310(R) (2011).
[22] S. Ryu and Y. Hatsugai, Phys. Rev. Lett. **89**, 077002 (2002).
[23] A.P. Schnyder and S. Ryu, Phys. Rev. B **84**, 060504(R) (2011); P.M.R. Brydon, A.P. Schnyder, and C. Timm Phys. Rev. B **84**, 020501(R) (2011).
[24] T.T. Heikkilä, N.B. Kopnin, and G.E. Volovik, Pis'ma v ZhETF **94**, 252 (2011).